TABLE D-10

FREQUENCY OF MATERNAL AND LATE MATERNAL DEATHS* BY YEAR MONTANA RESIDENTS. 2003-2007

| | 2003 | 2004 | 2005 | 2006 | 2007 |
|--|------|------|------|------|------|
| Total, all Maternal and Late Maternal Causes | 5 | 9 | 6 | 2 | 4 |
| Maternal causes (A34,O00-O95,O98-O99) | 2 | 2 | 1 | - | 2 |
| Pregnancy with abortive outcome (O00-O07) | - | - | - | - | - |
| Ectopic pregnancy (O00) | - | - | - | - | - |
| Spontaneous abortion (O03) | - | - | - | - | - |
| Medical abortion (O04) | - | - | - | - | - |
| Other abortion (O05) | - | ı | - | - | - |
| Other and unspecified pregnancy with abortive outcome (O01-O02,O06-O07) | - | ı | - | - | - |
| Other direct obstetric causes(A34,O10-O92) | - | 2 | - | - | 1 |
| Eclampsia and pre-eclampsia (O11,O13-O16) | - | - | - | - | 1 |
| Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72) | - | 1 | - | - | - |
| Complications predominately related to the puerperium (A34,O85-O92) | - | ı | - | - | - |
| Obstetrical tetanus (A34) | - | ı | - | - | - |
| Obstetric embolism (O88) | - | ı | - | - | - |
| Other complications predominately related to the puerperium O85-O87,O89-O92) | - | ı | - | - | - |
| All other direct obstetric causes(O10,O12,O21-O43,O47-O66,O68-O71,O73-O75) | 1 | 1 | - | - | - |
| Obstetric death of unspecified cause (O95) | - | | - | - | - |
| Indirect obstetric causes (O98-O99) | 2 | | 1 | - | 1 |
| Maternal causes more than 42 days after delivery or termination of pregnancy (O96-O97) | 3 | 7 | 5 | 2 | 2 |
| Death from any obstetric cause occurring more than 42 days but less than one year after delivery (O96) | 3 | 7 | 5 | 2 | 2 |
| Death from sequelae of direct obstetric causes (O97) | - | - | - | - | - |

^{*} A "MATERNAL DEATH" IS THE DEATH OF A FEMALE WHILE PREGNANT OR WITHIN 42 DAYS OF TERMINATION OF PREGNANCY, IRRESPECTIVE OF THE DURATION AND SITE OF PREGNANCY, FROM ANY CAUSE RELATED TO OR AGGREVATED BY THE PREGNANCY OR ITS MANAGEMENT, BUT NOT FROM ACCIDENT OR INCIDENTAL CAUSES.
MATERNAL CONDITIONS ARE THOSE ASSIGNED TO PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (ICD-10 CODES A34, 000-095, AND 098-099). "LATE MATERNAL DEATHS" ARE THOSE ATTRIBUTABLE TO THE SAME CAUSES BUT WHICH OCCURRED MORE THAN 42 DAYS AFTER TERMINATION OF THE PREGNANCY. THE 2003 REVISION OF THE MONTANA DEATH CERTIFICATE ADDED A PREGNANCY HISTORY ITEM WHICH INDICATES WHETHER THE DECEDENT WAS PREGNANT WITHIN THE 12 MONTHS PRIOR TO DEATH (ICD-10 CODES 096-097). THIS NEW ITEM MAKES THE IDENTIFICATION OF A GREATER NUMBER OF MATERNAL AND LATE MATERNAL DEATHS POSSIBLE.